

LETTER • OPEN ACCESS

A global map of mangrove forest soil carbon at 30 m spatial resolution

To cite this article: Jonathan Sanderman *et al* 2018 *Environ. Res. Lett.* **13** 055002

View the [article online](#) for updates and enhancements.

Related content

- [A global predictive model of carbon in mangrove soils](#)
Sunny L Jardine and Juha V Siikamäki
- [Estimating mangrove aboveground biomass from airborne LiDAR data: a case study from the Zambezi River delta](#)
Temilola Fatoyinbo, Emanuelle A Feliciano, David Lagomasino *et al.*
- [Measurement and monitoring needs, capabilities and potential for addressing reduced emissions from deforestation and forest degradation under REDD+](#)
Scott J Goetz, Matthew Hansen, Richard A Houghton *et al.*

Environmental Research Letters



LETTER

A global map of mangrove forest soil carbon at 30 m spatial resolution

OPEN ACCESS

RECEIVED

29 June 2017

REVISED

10 April 2018

ACCEPTED FOR PUBLICATION

13 April 2018



PUBLISHED

30 April 2018

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Jonathan Sanderman^{1,21} , Tomislav Hengl², Greg Fiske¹, Kylene Solvik¹, Maria Fernanda Adame³, Lisa Benson^{5,6}, Jacob J Bukoski⁷, Paul Carnell⁸, Miguel Cifuentes-Jara⁹, Daniel Donato¹⁹, Clare Duncan^{4,8}, Ebrahim M Eid^{10,20}, Philine zu Ermgassen^{17,18}, Carolyn J Ewers Lewis⁸, Peter I Macreadie⁸ , Leah Glass⁵, Selena Gress¹¹, Sunny L Jardine¹², Trevor G Jones^{5,13}, Eugène Ndemem Nsombo¹⁴, Md Mizanur Rahman¹⁵, Christian J Sanders¹⁶, Mark Spalding¹⁷ and Emily Landis¹⁷

¹ Woods Hole Research Center, 149 Woods Hole Road, Falmouth, MA 02540, United States of America

² ISRIC — World Soil Information, Wageningen, The Netherlands

³ Australian Rivers Institute, Griffith University, Nathan, QLD, Australia

⁴ Institute of Zoology, Zoological Society of London, Outer Circle, Regent's Park, London NW1 4RY, United Kingdom

⁵ Blue Ventures Conservation, London, United Kingdom

⁶ Centre for Environment, Fisheries and Aquaculture Science, Lowestoft, United Kingdom

⁷ Department of Environmental Science, Policy and Management, University of California, Berkeley, CA, United States of America

⁸ Centre for Integrative Ecology, School of Life and Environmental Sciences, Deakin University, Burwood, VIC, Australia

⁹ Forests, Biodiversity and Climate Change Program, CATIE 7170, Turrialba, Costa Rica

¹⁰ Botany Department, Faculty of Science, Kafr El-Sheikh University, Kafr El-Sheikh, Egypt

¹¹ School of Applied Sciences, Edinburgh Napier University, Edinburgh, Scotland

¹² School of Marine and Environmental Affairs, University of Washington, Seattle, WA, United States of America

¹³ Department of Forest Resources Management, University of British Columbia, Vancouver, BC, Canada

¹⁴ Institute of Fisheries and Aquatic Sciences, University of Douala, Douala, Cameroon

¹⁵ Graduate School of Agriculture, Kyoto University, Kyoto, Japan

¹⁶ National Marine Science Centre, Southern Cross University, Coffs Harbour, NSW, Australia

¹⁷ The Nature Conservancy, Arlington, VA, United States of America

¹⁸ School of Geosciences, University of Edinburgh, Edinburgh, United Kingdom

¹⁹ Washington State Department of Natural Resources, Olympia, WA United States of America

²⁰ Biology Department, College of Science, King Khalid University, Abha 61321, Saudi Arabia

²¹ Author to whom any correspondence should be addressed.

E-mail: jsanderman@whrc.org

Keywords: blue carbon, carbon sequestration, land use change, machine learning

Supplementary material for this article is available [online](#)

Abstract

With the growing recognition that effective action on climate change will require a combination of emissions reductions and carbon sequestration, protecting, enhancing and restoring natural carbon sinks have become political priorities. Mangrove forests are considered some of the most carbon-dense ecosystems in the world with most of the carbon stored in the soil. In order for mangrove forests to be included in climate mitigation efforts, knowledge of the spatial distribution of mangrove soil carbon stocks are critical. Current global estimates do not capture enough of the finer scale variability that would be required to inform local decisions on siting protection and restoration projects. To close this knowledge gap, we have compiled a large georeferenced database of mangrove soil carbon measurements and developed a novel machine-learning based statistical model of the distribution of carbon density using spatially comprehensive data at a 30 m resolution. This model, which included a prior estimate of soil carbon from the global SoilGrids 250 m model, was able to capture 63% of the vertical and horizontal variability in soil organic carbon density (RMSE of 10.9 kg m^{-3}). Of the local variables, total suspended sediment load and Landsat imagery were the most important variable explaining soil carbon density. Projecting this model across the global mangrove forest distribution for the year 2000 yielded an estimate of 6.4 Pg C for the top meter of soil with an $86\text{--}729 \text{ Mg C ha}^{-1}$ range across all pixels. By utilizing remotely-sensed mangrove forest cover change data, loss of soil carbon due to mangrove habitat loss between 2000 and 2015 was $30\text{--}122 \text{ Tg C}$ with $>75\%$ of this loss attributable to Indonesia, Malaysia and Myanmar. The resulting map products

from this work are intended to serve nations seeking to include mangrove habitats in payment-for-ecosystem services projects and in designing effective mangrove conservation strategies.

1. Introduction

Mangrove forests, occupying less than 14 million ha (Giri *et al* 2011), just 2.5% of the size of the Amazon rainforest, provide a broad array of ecosystem services (Barbier *et al* 2011). Mangroves are critical nursery habitats for fish, birds and marine mammals (Mumby *et al* 2004, Nagelkerken *et al* 2008), act as effective nutrient filters (Robertson and Phillips 1995), buffer coastal communities from storm surges (Gedan *et al* 2011) and support numerous rural economies (Spalding *et al* 2014, Temmerman *et al* 2013). These ecosystem service benefits have been valued at an average of 4200 US\$ ha⁻¹ yr⁻¹ in Southeast Asia (Brander *et al* 2012). Despite these ecosystem service benefits, mangroves are highly threatened by both urban expansion and other 'higher value' land uses because of their close proximity to major human settlements. There are no reliable estimates of original mangrove cover, but some authors have suggested that 35% or more of original cover may have been lost and wider areas have been degraded (Valiela *et al* 2001, Spalding *et al* 2010). Loss rates have slowed dramatically in the past 10–20 years in most areas, however they remain considerable, with rates up to 3.1% annually in some countries (Hamilton and Casey 2016). The major drivers of loss are conversion for aquaculture, especially shrimp farming, agriculture and urban development (Alongi 2002, Valiela *et al* 2001, Spalding *et al* 2010, Richards and Friess 2016) but loss due to extreme climatic events are also becoming more common (Duke *et al* 2017).

With the growing recognition that effective action on climate change will require a combination of emissions reductions and removals (Rockström *et al* 2017), protecting, enhancing and restoring natural carbon sinks have become political priorities (Boucher *et al* 2016, Grassi *et al* 2017). Mangrove forests can play an important role in carbon removals; in addition to being some of the most carbon-dense ecosystems in the world (Donato *et al* 2011), if kept undisturbed, mangrove forest soils act as long-term carbon sinks (Breithaupt *et al* 2012). As such, there is strong interest in developing policy tools to protect and restore mangroves through payment for ecosystem services (Friess *et al* 2016, Howard *et al* 2017).

Mangroves can store significant amounts of carbon in their biomass (Hutchison *et al* 2014); however, the vast majority of the ecosystem carbon storage is typically found in the soil (Donato *et al* 2011, Muddiyarso *et al* 2015, Sanders *et al* 2016). For example, Kauffman *et al* (2014) found that within the same estuary, soil carbon contributed 78% of total ecosystem

carbon storage in tall mangroves but 96%–99% of total ecosystem carbon in medium and low stature mangrove stands. Importantly, Kauffman *et al* (2014) found that conversion of these mangrove forests to shrimp ponds resulted in the loss of 90% of this carbon from the top 3 m of soil (612–1036 Mg C ha⁻¹). In addition to avoided emissions, many mangrove forest soils are accreting as sea level rises (Krauss *et al* 2014), providing continual carbon sequestration on the order of 1.3–2.0 Mg C ha⁻¹ yr⁻¹ (Breithaupt *et al* 2012, Chmura *et al* 2003). Clearly, there can be a major climate benefit to halting or even slowing the rate of mangrove conversion, with a rough potential estimated to be 25–122 Tg C yr⁻¹ (Pendleton *et al* 2012, Siikamäki *et al* 2012). For nations with large mangrove holdings, protection and restoration can make major contributions to meeting climate mitigation targets (Herr and Landis 2016).

While many mangrove forests do accumulate large quantities of soil carbon, others do not. There can be significant variability in soil carbon stocks across different mangrove forests (Jardine and Siikamäki 2014) but also within the same mangrove forest (Adame *et al* 2015, Kauffman *et al* 2011). Understanding the distribution of soil carbon in mangrove forests will be very important in prioritizing protection and restoration efforts for climate mitigation. The controls on soil carbon stocks are diverse and are likely scale dependent; however, some generalizations can be made. Mangrove forests, no matter how productive, will struggle to have high soil carbon stocks in the upper meter of soil if they receive large annual sediment loads. Mangrove forests in river deltas, such as the Sundarbans (Banerjee *et al* 2012) and the Zambezi river delta in Mozambique (Stringer *et al* 2016), typically only contain a few percent organic carbon throughout the soil profile. These locations may still have very high carbon stocks, but the density of carbon is low due to the high allocthonous input of mineral sediments. Conversely, forests with moderately low productivity can accumulate large amounts of soil carbon if they are in an isolated hydrogeomorphic setting (Ezcurra *et al* 2016). Within the same mangrove forest there are typically steep hydrogeomorphic gradients from the seaward to landward extent of the forest which results in zonation of both vegetation (Snedaker 1982) and soil carbon storage (Kauffman *et al* 2011, Ouyang *et al* 2017, Ewers Lewis *et al* 2018) but not necessarily for the same reasons. Within a similar hydrogeomorphic position, forest productivity and soil edaphic conditions (e.g. redox potential, pH, salinity) driving decomposition rates are often the dominant controls on soil carbon density. Consideration of this nested

hierarchy of controls will be necessary to successfully capture the variability in soil carbon at both local and global scales.

Accurate estimates and an understanding of the spatial distribution of mangrove soil carbon stocks are a critical first step in understanding climatic and anthropogenic impacts on mangrove carbon storage and in realizing the climate mitigation potential of these ecosystems through various policy mechanisms (Howard *et al* 2017). Previous global estimates (Atwood *et al* 2017, Jardine and Siikamäki 2014), do not capture enough of the finer scale spatial variability that would be required to inform local decisions on siting protection and restoration projects. To close this information gap, we have: (1) compiled and published a harmonized global database of the profile distribution of soil carbon under mangroves, (2) used this database to develop a novel machine-learning based data-driven statistical model of the distribution of carbon density using spatially comprehensive data at an ~ 30 m resolution, (3) projected the model results across global mangrove habitat for the year 2000 (Giri *et al* 2011), and (4) overlaid estimates of mangrove forest change between 2000 and 2012 (Hamilton and Casey 2016) to estimate potential soil carbon emissions from recent forest conversion.

2. Methods

2.1. Mangrove soil carbon database

A harmonized globally representative database (available at: [10.7910/DVN/OCYUIT](https://doi.org/10.7910/DVN/OCYUIT)) was compiled from peer-reviewed literature, grey literature and from contributions of unpublished data from a number of researchers and organizations. Details of database development and a statistical summary of the data are given in the supplemental information available at stacks.iop.org/ERL/13/055002/mmedia.

2.2. Spatial modelling of soil organic carbon

In order to maximize the utilization of available soil carbon data, we developed a machine learning-based model of organic carbon density (OCD) which models OCD as a function of depth (d), an initial estimate of the 0–200 cm organic carbon stock (OCS) from the global SoilGrids 250 m model (Hengl *et al* 2017), and a suite of spatially explicit covariate layers (X_p):

$$\text{OCD}(xyd) = d + \text{OCS}_{\text{SG}} + X_1(xy) + X_2(xy) + \dots X_p(xy)$$

where OCS_{SG} is the aggregated organic carbon stock estimated for 0–200 cm depth using global SoilGrids 250 m approach down-sampled from 250 m–30 m resolution, and xyd are the 3D coordinates northing easting and soil depth (measured to center of a horizon). Note here that we model spatial distribution

of OCD in three dimensions (soil depth used as a predictor) using all soil horizons layers at different depths, which means that a single statistical model can be used to predict OCD at any arbitrary depth. This 3D approach to modeling OCD reduces the need for making complex assumptions about the downcore trends in OCD, and maximizes the use of collected data.

The derived spatial prediction model is then used to predict OCD at standard depths 0, 30, 100, and 200 cm, so that the organic carbon stock (OCS) can be derived as a cumulative sum of the layers down to the prediction depth for every 30 m pixel identified as having mangrove forest in the year 2000 (Giri *et al* 2011). Importantly, we found that there is a spatial mismatch between the global mangrove forest distribution (GMFD) of Giri *et al* (2011) and satellite imagery (figure S2). To best resolve this spatial mismatch, we have adjusted the GMFD by growing all vectors by one pixel (~ 30 m) and then filtering out any pixel that falls over water by using Landsat NIR band (see SI for more details).

Environmental covariates have been compiled to represent the postulated major controls on OCS in soils generally (McBratney *et al* 2003) and specifically for mangrove ecosystems (Balke and Friess 2016). Covariates included:

1. Vegetation characteristics including percent forest cover (Hansen *et al* 2013) and Landsat bands 3 (red), 4 (near infrared), 5 (shortwave infrared) and 7 (shortwave infrared) for the year 2000 (Hanson *et al* 2013) retrieved from http://earthenginepartners.appspot.com/science-2013-global-forest/download_v1.3.html;
2. Digital elevation data, which at or near sea-level approximately follows forest canopy height (Simard *et al* 2006), from the shuttle radar topography mission (SRTM GL1; NASA, 2013) was retrieved from <https://lpdaac.usgs.gov>, maintained by the NASA EOSDIS Land Processes Distributed Active Archive Center (LP DAAC) at the USGS/Earth Resources Observation and Science (EROS) Center, Sioux Falls, South Dakota;
3. Long-term averaged (1990–2010) monthly sea surface temperature (SST) averaged into four seasons were generated in Google Earth Engine from NOAA AVHRR Pathfinder Version 5.2 Level 3 Collated data (Casey *et al* 2010) and downsampled to 30 m resolution using bicubic resampling;
4. The M2 tidal elevation amplitude product (FES2012) from a global hydrodynamic tidal model which assimilates altimetry data from multiple platforms was used to represent tidal range at each location. The FES2012 product was produced by Noveltis, Legos and CLS Space Oceanography Division and distributed by Aviso, with support from Cnes (www.aviso.altimetry.fr/);

5. Averaged (2003–2011) monthly total suspended matter (TSM) averaged into four seasons estimated from MERIS imagery collected by the European Space Agency's Envisat satellite. Processed and validated TSM data was retrieved from the GlobColour project (<http://hermes.acri.fr>).
6. A mangrove typology map delineating mangroves into estuaries and then either organogenic or mineralogenic based on an analysis of TSM and tidal amplitude data (Zu Ermgassen, *unpublished data*).

Sea surface temperature, tidal amplitude and TSM are 4 km resolution ocean products and needed to be extrapolated to each pixel containing mangrove forest. Missing values in the sea surface temperature, tidal amplitude and TSM were first filled-in using spline interpolation in SAGA GIS, then down-scaled to 30 m resolution using bicubic resampling in GDAL. Including SoilGrids and depth, there were a total of 20 covariates used in building the mangrove OCD model.

The ability of the training points to represent the entire covariate space of the global mangrove domain was assessed by conducting a principal components analysis (PCA) on 15 000 randomly selected points and the 1613 points used in the spatial model. Spatial variables were detrended and centered by subtracting the mean and dividing by the standard deviation (s.d.) before entering into the PCA analysis.

Soil carbon typically varies in highly non-linear ways with depth and across the landscape and as such the ability of standard parametric models to capture this variation is limited (Jardine and Siikamäki 2014, Hengl *et al* 2017). Here we model the spatial (xyd) distribution of OCD using a machine learning random forest model implemented in the ranger package (Wright and Ziegler 2015) in the R environment for statistical computing (R Core Team 2000). Given the clustered nature of the point data, we have implemented a spatially balanced random forest model design. Model performance was assessed with a 5 fold (Leave-Location-Out) cross-validation procedure where 20% of complete locations were withheld in each model refitting (Gasch *et al* 2015). The relative importance of using SoilGrids as a covariate was assessed by implementing the cross-validation procedure with and without this variable.

Finally, prediction error of OCS for 0–1 m depth was derived for ± 1 s.d. based on the quantile regression approach of Meinshausen (2006) and implemented in R via the ranger package. This procedure is relatively computationally demanding so a random subset of approximately 15 000 points were selected to calculate prediction errors. All modeling was run on ISRIC High Performance Computing servers with 48 cores of 256 GB RAM.

2.3. Data analysis


Soil carbon stocks were calculated for the global extent of mangroves for the year 2000 by summing the OCS in each pixel for 1 and 2 m depths. Country level carbon stocks were also calculated for the same depths. Given the fringing nature of mangroves, a global spatial vector data layer was built that allocated the offshore area for each country where mangrove forests can be found. It was derived from the Exclusive Economic Zone for each country. This layer was then dissolved with the onshore areas for each associated country and subsequently used to quantify mangrove OCS tonnage and areal extent.

Potential loss of OCS due to mangrove habitat conversion was calculated between 2000 and 2015 by summing the OCS in mangrove forest pixels which were identified to be deforested. While this analysis cannot distinguish between natural and anthropogenic disturbance, human-driven land use change is believed to be by far the dominant driver of deforestation in mangrove ecosystems (Alongi 2002, Murdiyarso *et al* 2015). We define deforested using the Global Forest Change dataset (Hansen *et al* 2013) available online from: <http://earthenginepartners.appspot.com/science-2013-global-forest>. We chose to use this approach for estimating deforestation instead of using the derived mangrove tree cover loss data produced by Hamilton and Casey (2016) as used by Atwood *et al* (2017) because the Hamilton and Casey (2016) analysis only considered forested area as area actually covered by trees (i.e. if a 100 ha forest has 80% tree cover then it is counted as 80 ha of forest). In our opinion, this definition mischaracterizes forest area extent. Next, an estimate of the soil carbon emissions associated with land use conversion is needed. The amount of OCS lost can be highly variable and depends on the new land use (Kauffman *et al* 2014, 2016b, Jones *et al* 2015) and probably on soil properties. Pendleton *et al* (2012) used a 25%–100% loss range. Donato *et al* (2011) used a low estimate of 25% of the OCS in top 30 cm and 75% in top 30 cm + 35% from deeper layers as a high estimate. Expanding on earlier work, Kauffman *et al* (2017) found that on average 54% of belowground carbon (soil + roots) to 3 m was lost after conversion to shrimp ponds and pastures. Given the limited number of studies comparing soil OCS change with land use change, in this work we adopt the same 25%–100% range as used by Pendleton *et al* (2012) applied to the first meter of soil. Finally, country level statistics for OCS loss were calculated as described above. All global and country level analyses were performed on the 30 m resolution dataset in Google Earth Engine (Gorelick *et al* 2017).

3. Results

3.1. Model results

The random forest model was successful in capturing the major variation in OCD across the mangrove



database (figure 1(a)) with an R^2 of 0.84 and a root mean square error (RMSE) of 6.9 kg m^{-3} compared to the mean OCD value of 29.6 kg m^{-3} . Cross-validation results yield an R^2 of 0.63 and an RMSE of 10.9 kg m^{-3} (figure 1(b)), which is the de-facto mapping accuracy to be expected on the field. Low OCD values were slightly over-predicted and high OCD values were under-predicted (figure 1(b)). The initial OCS prediction from SoilGrids 250 m was the most important variable explaining mangrove OCD. Running the 5 fold cross-validation without and with SoilGrids indicated that this single variable explained improved model performance by nearly 50% (R^2 increased from 0.42–0.63). Seasonal total suspended matter (TSM), depth of sample, mangrove tree cover, Landsat Red band, sea surface temperature (SST), and tidal range were the next ten most important variables, respectively (figure 1(c)). Quantile regression analysis indicated that the full uncertainty (± 1 s.d.) about a mean prediction of carbon stocks to 1 m depth averaged 40.4% of the mean OCS with lowest relative uncertainty in the most carbon-rich mangrove forests (figure S7).

3.2. Mangrove soil carbon storage

Projection of the mangrove OCD model to global mangrove forests revealed the distribution of soil carbon storage in these ecosystems (figure 2). The mean (± 1 s.d.) OCS to 1 m depth was $361 \pm 136 \text{ Mg C ha}^{-1}$ with a range of $86\text{--}729 \text{ Mg C ha}^{-1}$. At the national level (table S1), Bangladesh had the lowest per ha stocks, averaging just 127 Mg C ha^{-1} followed by China and the nations bordering the Persian Gulf and Red Sea with an average OCS of 214 and 233 Mg C ha^{-1} , respectively. The highest per ha stocks were found in many of the Pacific island nations, averaging 505 Mg C ha^{-1} with much of Southeast Asia ranking well above the global mean.

While the national level comparisons are revealing, by modeling at a 30 m resolution much richer details of potential within forest variation in OCS are

seen (figure 2). Mangrove forests dominated by sediment laden fluvial inputs typically have consistently low OCS as seen in the Sundarbans and Madagascar (figures 2(a) and (e)). In non-deltaic mangroves, the model appears to have captured the large zonal variation in OCS that is often observed in field studies (figures 2(b) and (c)).


3.3. Soil carbon loss due to habitat loss

Utilizing the Hanson *et al* (2013) global deforestation analysis (figure S8), we found that 278049 ha (1.67% of total) of area identified as mangrove habitat in the year 2000 was deforested resulting in the committed emission of $30.4\text{--}122 \text{ Tg C}$ ($111\text{--}447 \text{ Tg CO}_2$) from mangrove forest soils due to land use change between 2000 and 2015 (figure 3). The relative rank of nations in terms of loss of mangrove forest area and OCS were often not the same (table S1). Indonesia alone was responsible for 52% of this global loss with Malaysia and Myanmar representing another 25% of the global total loss (figure 3(c)). When visualized as a percent loss from year 2000 stocks, a slightly different pattern emerged (figure 3(d)). Guatemala had the highest percent loss of mangrove OCS (0.9%–6.8%) followed by several southeast Asian nations, but high percent losses were also found in several Caribbean island nations as well as the United States and several west African countries.

4. Discussion

4.1. Amount and distribution of Mangrove SOC

Our new estimate of global mangrove OCS of 6.4 Pg C in the upper meter and 12.6 Pg C to 2 m is largely consistent with past efforts to calculate this value (Donato *et al* 2011, Jardine and Siikamäki 2014, Sanders *et al* 2016). However, our estimate is double that of Atwood *et al* (2017) primarily due to their use of the Hamilton and Casey (2016) estimate of mangrove extent instead of Giri *et al* (2011). Importantly, by using an environmental covariate model,




we have been able to make plausible estimates for regions where no sampling has taken place instead of relying on global mean values (i.e. Atwood *et al* 2017).

The total amount of soil carbon was similar in our analysis and the most comparable analysis, that of Jardine and Siikamäki (2014), but the spatial distribution of carbon-rich versus carbon-poor mangroves varied substantially. For example, we found much higher OCS levels in West Africa than in East African nations (figure 2 and table S1) but the reverse was found by Jardine and Siikamäki (2014). Large discrepancies were also found for Colombia, Sri Lanka and many of the countries bordering the Red Sea. These differences were most likely driven by lack of data in those regions at the time of the analysis by Jardine and Siikamäki (2014) given that nearly half the data in our database was collected after their study was published. Additionally, our analysis suggested a much larger range in OCS ($86\text{--}729\text{ Mg C ha}^{-1}$) compared to $272\text{--}703\text{ Mg C ha}^{-1}$ in the analysis of

Jardine and Siikamäki (2014). This difference in range was likely due to the inclusion of more data from sub-tropical and temperate mangroves (figure S3).

The depth trend analysis (figure S6) and random forest variable importance (figure 1(b)) both indicated that depth should be considered in calculation of OCS. For locations that were either stable peat domes or located in estuaries receiving large annual sediment loads, a stable OCD profile distribution would be expected and this was found for many sites (figures S6(a) and (e)). However, where mangroves are growing in a mineral matrix that is receiving only low sediment loads, a decline in OCD may be expected as the carbon inputs from the productive mangrove forest would be concentrated in the surface horizons (figure S6(b)). Still in other cases (figure S6(c)), changes in hydrologic/sediment regimes can lead to irregular depth patterns or even an increase in OCD with depth.



While total area of mangroves was a key determinant of total soil carbon storage, amongst the top 25 mangrove OCS holding nations, there was a nearly even split between nations with smaller area of high soil carbon density forests and those nations with lots of low soil carbon density forests (figure 4). Indonesia was the clear exception to this trend with the largest mangrove holdings which also contain rich carbon stocks resulting in Indonesia alone holding nearly 25% of the world's mangrove OCS.

Compared to terrestrial carbon pools, mangrove forests rank low due to their limited spatial extent. For example in the upper meter of soil, permafrost affected soils are estimated to store 472 ± 27 Pg C (Hugelius *et al* 2014), tropical forests contain ~ 188 Pg C, and soils under permanent cropping contain ~ 150 Pg C (table 1). However, on an equal area basis, mangrove forests on average store more soil carbon than most other ecosystems (table 1). Importantly, our

analysis has demonstrated mangrove soil carbon is highly variable and many mangroves actually store fairly modest amounts of carbon in the upper one or two meters of soil. While not the focus of this analysis, it is important to point out that while some mangrove forests store modest levels of OCS in the upper meter of soil, they can have high sequestration rates and conversely carbon-dense mangroves can have low annual sequestration rates (Lovell *et al* 2010, MacKenzie *et al* 2016).

4.2. Drivers of soil carbon storage

Our spatial modelling framework, in which global predictions were combined with local high resolution images, was successful as the general patterns of carbon variation from SoilGrids 250 m were maintained, while the spatial detail was significantly improved by moving from 250 m–30 m spatial resolution. The initial SoilGrids 250 m OCS prediction (Hengl *et al* 2017)

Table 1. Soil organic carbon stocks (mean with 5th–95th percentile in parentheses) and total storage for different terrestrial ecosystems compared to mangrove forests.

Land cover category (IGBP code) ^a	Area (10 ⁶ ha)	1 m soil organic carbon stock	
		Pg C	Mg C ha ⁻¹
Mangrove forest ^b	16.6	6.4	361 (94–628)
Gelisols (permafrost soils) ^c	1878	472.0	389 (178–691)
Evergreen Needleleaf forest (1) ^d	286	60.0	210 (121–346)
Evergreen Broadleaf forest (2)	1248	188.4	151 (85–271)
Deciduous Needleleaf forest (3)	116	29.3	253 (163–412)
Deciduous Broadleaf forest (4)	165	22.1	134 (83–223)
Mixed forest (5)	771	152.8	198 (93–343)
Closed shrublands (6)	56	6.2	110 (39–223)
Open shrublands (7)	1933	325.8	169 (49–329)
Woody savannas (8)	1179	185.8	158 (82–274)
Savannas (9)	1010	112.9	112 (52–201)
Grasslands (10)	1810	280.1	155 (56–289)
Permanent wetlands (11) ^e	104	25.1	241 (114–474)
Croplands (12)	1177	149.6	127 (60–200)
Cropland/Natural veg. mosaic (14)	868	117.7	136 (58–238)


^a data for MODIS-based IGBP land cover classes (Friedl *et al* 2010) extracted from 1 m OCS map for the year 2010 produced by Sanderman *et al* (2017).

^b mangrove area and OCS data from this study.

^c permafrost area from Tarnocai *et al* (Tarnocai *et al* 2009), OCS data from Hugelius *et al* (2014).

^d some overlap between class 1 (evergreen Needleleaf forest) and gelisols.

^e class 11 (permanent wetlands) likely has overlap with mangrove area.



was based upon machine learning algorithms using 237 covariates that covered the main state factors of soil formation (Jenny 1994)—climate, relief, living organisms (vegetation) and parent material—and was four times more important in predicting mangrove OCD than any of the local covariates. However, the local covariates allowed for a much more refined picture of the spatial variation in OCS within mangrove forests (figure 2) which was not captured in the 250 m resolution SoilGrids 250 m prediction. Covariates related to hydrogeomorphology (TSM and tidal range), as hypothesized, were important predictors of local variation in OCD. Both TSM and tidal range were strongly negatively correlated with OCD suggesting that locations with either high sediment loads or strong tidal flushing do not accumulate large carbon


stocks. The mangrove typology ended up being uninformative likely because most of the data that went into this classification was already captured in the model.

Covariates related to mangrove biomass (SRTM elevation and Landsat bands) were also important in explaining the local distribution of OCD (figure 1(b)). However, as pointed out by Bukoski *et al* (2017), it is unclear whether the importance of these vegetation-related data are causal drivers of differences in OCD or they just happen to co-vary in the same way as OCD. To further explore the relationship between forest biomass carbon and soil carbon storage, we extracted aboveground biomass data from Hutchison *et al* (2014) and compared it to our OCS results (figure 5). While a clear positive trend was found between biomass and soil carbon storage ($R^2 = 0.26$), there is clearly a lot of variance especially at lower biomass levels where nearly the full range in OCS can be found.

Depth below the soil surface was an important covariate in modeling OCD distribution (figure 1(b)). This finding was supported by the database depth trend analysis (figure S6) which indicated that a flat depth distribution of OCD would be an incorrect assumption 37%–64% of the time. These findings suggest that the simple scaling performed in figure S5 and in several previous assessments (Bukoski *et al* 2017, Jardine and Siikamäki 2014, Atwood *et al* 2017) may not be an accurate estimation of total OCS especially when extrapolating from a surface horizon sample alone.

4.3 Soil carbon loss due to land conversion (2000–2015)

Our analysis suggests that mangrove soils have lost or are at least committed to losing 30.4–122 Tg C due to the land use conversion that occurred between the



years 2000 and 2015 (figure 3). Given that at the global level the rate of mangrove forest lost was consistent over this time period (Hamilton and Casey 2016), we estimated an annual soil carbon emission of 2.0–8.1 Tg C yr⁻¹. This value is significantly lower than previous estimates (Donato *et al* 2011, Pendleton *et al* 2012) for two reasons. First, we use remote sensing-based measurements of actual mangrove loss instead of applying a large range of annual conversion rates, which are notoriously variable according to their source (Friess and Webb 2011). Second, we have summed the actual OCS values for each of the pixels where land conversion has taken place (i.e. figure S8) instead of applying a conversion rate to a mean OCS value.

The three nations of Indonesia, Malaysia and Myanmar contributed 77% of global mangrove OCS loss for this time period (figure 3). Despite similar area loss (figure 3(a)), Malaysia lost approximately twice as much soil carbon as Myanmar due to the large differences in carbon density between these two nations (mean OCS = 485 ± 57 versus 245 ± 63 Mg C ha⁻¹, respectively). This comparison highlights the importance of using local OCS values for estimating carbon emissions attributed to mangrove conversion.

Not all land use conversions result in equal loss of OCS. Conversion of mangrove forest to shrimp ponds results in a rapid and near complete loss of carbon in the upper meter of soil (Kauffman *et al* 2014), as well as losses deeper in the soil profile (Kauffman *et al* 2017). Conversion to other agricultural uses such as pasture for beef production (Kauffman *et al* 2016b) and cereal crops (Andreetta *et al* 2016) also appear to result in large soil carbon emissions. However, mangrove degradation and loss due to over harvesting for fuelwood (Jones *et al* 2015) or due to natural

disturbance (Cahoon *et al* 2003) likely leads to more moderate emissions as decomposition and erosion exceed new plant carbon inputs.

It is important to note that nearly all available data on OCS loss due to conversion to other land uses come from organogenic mangrove forests. In a mineral-dominated mangrove systems with only a few percent sediment OCC, we would not expect the same level of carbon loss as when peat deposits are drained or removed. In fact, reclamation of deltaic sediments for paddy rice cultivation can lead to increases in OCS (Kalbitz *et al* 2013), although methane emissions would be expected to increase. Additionally, if mangrove habitat is lost due to deforestation without a change in hydrologic regime, mineral-dominated mangroves can continue to accrete carbon, but at a lower rate than in a system that has additional organic matter inputs from the mangroves themselves (Pérez *et al* 2017).

4.4. Limitations and uncertainties

While we endeavored to ensure that the model input data was of the highest quality possible, there undoubtedly remain unknown errors in the database which are contributing to model error. Machine learning models are particularly sensitive to outlier values and extrapolation (Murphy 2012). Various research groups use different methods for determining the organic carbon concentration (OCC) of a sample with not all publications reporting whether or not results were corrected for occurrence of inorganic carbon or whether or not roots were excluded before further processing. Bulk density (BD) is a difficult parameter to measure accurately in many soils, and based on our analysis of BD versus OCC (figure S1) some reported data are unlikely to be accurate. We

developed a procedure to correct potential BD errors, but a pedotransfer function gives only an approximation of the true value. Given the importance of depth in our models (figure 1(b)), it was unfortunate that so many investigations only report OCS for large depth increments. We suggest that future studies using the common practice of collecting subsamples within larger horizon increments (e.g. Kauffman and Donato 2012, Kauffman *et al* 2016a) report the specific depth increment of the sample rather than that of the entire core. This additional level of transparency in the data would allow mass-preserving splines (Bishop *et al* 1999) to be fit through the distinct measurement intervals, resulting in unbiased estimates of OCS.

The largest uncertainty in the input data likely resulted from imperfect information on plot location. Whether, accidental or purposeful (i.e. not wanting to identify exact locations), spatially misplaced data in publications are of limited utility in geospatial applications. All of the covariate data for each mangrove point were selected from spatial layers resulting in the potential for a mismatch between the recorded OCS and the spatial predictors. In this study, we visually inspected all coordinates against Google Earth imagery and our adjusted mangrove domain, and then contacted many authors to seek further information and manually adjusted coordinates when we were confident that the adjustments lead to better spatial location. In the final analysis, 199 soil profiles had to be excluded from analysis because we could not confidently locate these plots within the adjusted mangrove spatial domain.

5. Conclusions

This work has produced three resources which we hope to be of significant value to the blue carbon research and management communities: 1) a large harmonized database of soil carbon data from mangrove ecosystems; 2) high-resolution (30 m) predictions with error of soil carbon stocks across all mangrove forests globally; 3) estimates of potential soil carbon losses due to mangrove habitat loss between 2000 and 2015. By using a statistical data-driven model, we have been able to produce credible estimates OCS for the numerous mangrove regions where no field data exist. We found that mangrove OCS is highly variable (86–729 Mg C ha⁻¹ in the top meter) but that much of the variability could be captured using spatially-comprehensive predictors in a machine-learning framework. Of the 6400 Tg C in the upper meter of soil, 30–122 Tg have likely been lost due to deforestation since the year 2000 with 77% of this loss attributed to Indonesia, Malaysia and Myanmar. These spatially-explicit estimates of mangrove soil carbon storage and loss will provide a practical first step for enabling nations to prioritize mangrove protection as part of their climate mitigation and adaptation plans.

Acknowledgments

We would like to thank A Andreetta, M Osland, J M Smoak, A DelVecchia, M E Gonnee, R K Bhomia and J Kelleway for providing additional data from their publications. S-T Kang for translating and extracting data from Chinese language papers. R K Bhomia acknowledges CIFOR SWAMP project, USFS International Program and USAID for funding. M Rahman acknowledges support from USAID for funding. PM and CJS acknowledge support from the Australian Research Council (DE130101084 and LP160100242) and (DE160100443 and DP150103286), respectively. CD acknowledges support from the Rufford Foundation and Darwin Initiative for funding. JS, TH, GF and KS were supported by an anonymous gift to The Nature Conservancy. MFA was supported by funding from the Queensland Government through the Advance Queensland Fellowship. ISRIC is a non-profit organization primarily funded by the Dutch government.

Data availability

The mangrove soil carbon database and model outputs can be downloaded from Harvard dataverse at <https://dataverse.harvard.edu/dataset.xhtml?persistentId=doi:10.7910/DVN/OCYUIT>.

ORCID iDs

Jonathan Sanderman  <https://orcid.org/0000-0002-3215-1706>

Peter I Macreadie  <https://orcid.org/0000-0001-7362-0882>

References

- Adame M F, Santini N S, Tovilla C, Vázquez-Lule A, Castro L and Guevara M 2015 Carbon stocks and soil sequestration rates of tropical riverine wetlands *Biogeosciences* **12** 3805–18
- Alongi D M 2002 Present state and future of the world's mangrove forests *Environ. Conserv.* **29** 331–49
- Andreetta A, Huertas A D, Lotti M and Cerise S 2016 Land use changes affecting soil organic carbon storage along a mangrove swamp rice chronosequence in the Cacheu and Oio regions (northern Guinea-Bissau) *Agric. Ecosyst. Environ.* **216** 314–21
- Atwood T B *et al* 2017 Global patterns in mangrove soil carbon stocks and losses *Nat. Clim. Change* **7** 523–8
- Balke T and Friess D A 2016 Geomorphic knowledge for mangrove restoration: a pan-tropical categorization *Earth Surf. Process. Landforms* **41** 231–9
- Banerjee K, Chowdhury M R, Sengupta K, Sett S and Mitra A 2012 Influence of anthropogenic and natural factors on the mangrove soil of Indian Sundarbans wetland *Arch. Environ. Sci.* **6** 80–91
- Barbier E B, Hacker S D, Kennedy C, Koch E W, Stier A C and Silliman B R 2011 The value of estuarine and coastal ecosystem services *Ecol. Monogr.* **81** 169–93
- Bishop T F A, McBratney A B and Laslett G M 1999 Modelling soil attribute depth functions with equal-area quadratic smoothing splines *Geoderma* **91** 27–45

- Boucher O, Bellassen V, Benveniste H, Ciais P, Criqui P, Guivarch C, Le Treut H, Mathy S and Séférian R 2016 Opinion: in the wake of Paris Agreement, scientists must embrace new directions for climate change research *Proc. Natl Acad. Sci. USA* **113** 7287–90
- Brander L, Wagtendonk A, Hussain S, McVittie A, Verburg P H, de Groot R S and van der Poeg S 2012 Ecosystem service values for mangroves in Southeast Asia: a meta-analysis and value transfer application *Ecosyst. Serv.* **1** 62–9
- Breithaupt J L, Smoak J M, Smith T J, Sanders C J and Hoare A 2012 Organic carbon burial rates in mangrove sediments: strengthening the global budget *Glob. Biogeochem. Cycles* **26** GB3011
- Bukoski J J, Broadhead J S, Donato D C, Murdiyarso D and Gregoire T G 2017 The use of mixed effects models for obtaining low cost ecosystem carbon stock estimates in mangroves of the Asia Pacific *PLoS ONE* **12** e0169096
- Cahoon D R, Hengstler P, Rybczyk J, McKee K L, Proffitt C E and Perez B C 2003 Mass tree mortality leads to mangrove peat collapse at Bay Islands, Honduras after Hurricane Mitch *J. Ecol.* **91** 1093–105
- Casey K S, Brandon T B, Cornillon P and Evans R 2010 The past, present, and future of the AVHRR Pathfinder SST program *Oceanography from Space* (Dordrecht: Springer Netherlands) pp 273–87
- Chmura G L, Anisfeld S C, Cahoon D R and Lynch J C 2003 Global carbon sequestration in tidal, saline wetland soils *Glob. Biogeochem. Cycles* **17** 1111
- Donato D C, Kauffman J B, Murdiyarso D, Kurnianto S, Stidham M and Kanninen M 2011 Mangroves among the most carbon-rich forests in the tropics *Nat. Geosci.* **4** 293–7
- Duke N C, Kovacs J M, Griffiths A D, Preece L, Hill D J E, Van Oosterzee P, Mackenzie J, Morning H S and Burrows D 2017 Large-scale dieback of mangroves in Australia's Gulf of Carpentaria: a severe ecosystem response, coincidental with an unusually extreme weather event *Mar. Freshw. Res.* **68** 1816–29
- Ewers Lewis C J, Sanderman P E, Baldock J and Macreadie P I 2018 Variability and vulnerability of coastal blue carbon stocks: a case study from Southeast Australia *Ecosystems* **21** 263–79
- Ezcurra P, Ezcurra E, Garcillán P P, Costa M T and Aburto-Oropeza O 2016 Coastal landforms and accumulation of mangrove peat increase carbon sequestration and storage *Proc. Natl Acad. Sci.* **113** 4404–9
- Friedl M A, Sulla-Menashe D, Tan B, Schneider A, Ramankutty N, Sibley A and Huang X 2010 MODIS collection 5 global land cover: algorithm refinements and characterization of new datasets *Remote Sens. Environ.* **114** 168–82
- Friess D A, Thompson B S, Brown B, Amir A A, Cameron C, Koldewey H J, Sasmito S D and Sidik F 2016 Policy challenges and approaches for the conservation of mangrove forests in Southeast Asia *Conserv. Biol.* **30** 933–49
- Friess D A and Webb E L 2011 Bad data equals bad policy: how to trust estimates of ecosystem loss when there is so much uncertainty? *Environ. Conserv.* **38** 1–5
- Gasch C K, Hengl T, Gräler B, Meyer H, Magney T S and Brown D J 2015 Spatio-temporal interpolation of soil water, temperature, and electrical conductivity in 3D+ T: the cook agronomy farm data set *Spat. Stat.* **14** 70–90
- Gedan K B, Kirwan M L, Wolanski E, Barbier E B and Silliman B R 2011 The present and future role of coastal wetland vegetation in protecting shorelines: answering recent challenges to the paradigm *Clim. Change* **106** 7–29
- Giri C, Ochieng E, Tieszen L L, Zhu Z, Singh A, Loveland T, Masek J and Duke N 2011 Status and distribution of mangrove forests of the world using earth observation satellite data *Glob. Ecol. Biogeogr.* **20** 154–9
- Gorelick N, Hancher M, Dixon M, Ilyushchenko S, Thau D and Moore R 2017 Google Earth engine: planetary-scale geospatial analysis for everyone *Remote Sens. Environ.* **202** 18–27
- Grassi G, House J, Dentener F, Federici S, den Elzen M and Penman J 2017 The key role of forests in meeting climate targets requires science for credible mitigation *Nat. Clim. Change* **7** 220–6
- Hamilton S E and Casey D 2016 Creation of a high spatio-temporal resolution global database of continuous mangrove forest cover for the 21st century (CGMFC-21) *Glob. Ecol. Biogeogr.* **25** 729–38
- Hansen M C *et al* 2013 High-resolution global maps of 21st century forest cover change *Science* **342** 850–3
- Hengl T *et al* 2017 SoilGrids 250 m: Global gridded soil information based on machine learning *PLoS ONE* **12** e0169748
- Herr D and Landis E 2016 *Coastal blue carbon ecosystems. Opportunities for Nationally Determined Contributions. Policy Brief* (Gland: IUCN and Washington, DC: TNC)
- Howard J, Sutton-Grier A, Herr D, Kleypas J, Landis E, Mcleod E, Pidgeon E and Simpson S 2017 Clarifying the role of coastal and marine systems in climate mitigation *Front. Ecol. Environ.* **15** 42–50
- Hugelius G *et al* 2014 Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps *Biogeosciences* **11** 6573–93
- Hutchison J, Manica A, Swetnam R, Balmford A and Spalding M 2014 Predicting global patterns in mangrove forest biomass *Conserv. Lett.* **7** 233–40
- Jardine S L and Siikamäki J V 2014 A global predictive model of carbon in mangrove soils *Environ. Res. Lett.* **9** 104013
- Jenny H 1994 *Factors of soil formation: a system of quantitative pedology* (Courier Corporation)
- Jones T *et al* 2015 The dynamics, ecological variability and estimated carbon stocks of mangroves in Mahajamba Bay, Madagascar *J. Mar. Sci. Eng.* **3** 793–820
- Kalbitz K *et al* 2013 The carbon count of 2000 years of rice cultivation *Glob. Change Biol.* **19** 1107–13
- Kauffman J B, Arifanti V B, Basuki I, Kurnianto S, Novita N, Murdiyarso D, Donato D C and Warren M W 2016a Protocols for the measurement, monitoring, and reporting of structure, biomass, carbon stocks and greenhouse gas emissions in tropical peat swamp forests *Working paper 221* (Bogor: CIFOR)
- Kauffman J B, Hernandez Trejo H, del Carmen Jesus Garcia M, Heider C and Contreras W M 2016b Carbon stocks of mangroves and losses arising from their conversion to cattle pastures in the Pantanos de Centla, Mexico *Wetl. Ecol. Manage.* **24** 203–16
- Kauffman J B, Arifanti V, Trejo H, del Carmen Jesús García M, Norfolk J, Hadriyanto D, Cifuentes-Jara M, Murdiyarso D and Cifuentes-Jara M 2017 The jumbo carbon footprint of a shrimp: carbon losses from mangrove deforestation *Front. Ecol. Environ.* **15** 183–8
- Kauffman J B and Donato D 2012 Protocols for the measurement, monitoring and reporting of structure, biomass and carbon stocks in mangrove forests *Working paper 86* (Bogor: CIFOR)
- Kauffman J B, Heider C, Cole T G, Dwire K A and Donato D C 2011 Ecosystem carbon stocks of micronesia mangrove forests *Wetlands* **31** 343–52
- Kauffman J B, Heider C, Norfolk J and Payton F 2014 Carbon stocks of intact mangroves and carbon emissions arising from their conversion in the Dominican Republic *Ecol. Appl.* **24** 518–27
- Krauss K W, McKee K L, Lovelock C E, Cahoon D R, Saintilan N, Reef R and Chen L 2014 How mangrove forests adjust to rising sea level *New Phytol.* **202** 19–34
- Lovelock C E, Sorrell B K, Hancock N, Hua Q and Swales A 2010 Mangrove forest and soil development on a rapidly accreting shore in New Zealand *Ecosystems* **13** 437–51
- MacKenzie R A, Foulk P B, Klump J V, Weckerly K, Purbospito J, Murdiyarso D, Donato D C and Nam V N 2016 Sedimentation and belowground carbon accumulation rates in mangrove forests that differ in diversity and land use: a tale of two mangroves *Wetl. Ecol. Manage.* **24** 245–61

- McBratney A, Mendonça Santos M and Minasny B 2003 On digital soil mapping *Geoderma* **117** 3–52
- Meinshausen N 2006 Quantile regression forests *J. Mach. Learn. Res.* **7** 983–99
- Mumby P J *et al* 2004 Mangroves enhance the biomass of coral reef fish communities in the Caribbean *Nature* **427** 533–6
- Murdiyarmo D, Purbopuspito J, Kauffman J B, Warren M W, Sasmito S D, Donato D C, Manuri S, Krisnawati H, Taberima S and Kurnianto S 2015 The potential of Indonesian mangrove forests for global climate change mitigation *Nat. Clim. Change* **5** 8–11
- Murphy K P 2012 *Machine Learning: A Probabilistic Perspective* (Cambridge, MA: MIT)
- Nagelkerken I *et al* 2008 The habitat function of mangroves for terrestrial and marine fauna: a review *Aquat. Bot.* **89** 155–85
- Ouyang X, Lee S Y and Connolly R M 2017 Structural equation modelling reveals factors regulating surface sediment organic carbon content and CO₂ efflux in a subtropical mangrove *Sci. Total Environ.* **578** 513–22
- Pendleton L *et al* 2012 Estimating global ‘blue carbon’ emissions from conversion and degradation of vegetated coastal ecosystems *PLoS ONE* **7** e43542
- Pérez A, Machado W, Gutierrez D, Stokes D, Sanders L, Smoak J M, Santos I and Sanders C J 2017 Changes in organic carbon accumulation driven by mangrove expansion and deforestation in a New Zealand estuary *Estuar. Coast. Shelf Sci.* **192** 108–16
- Richards D R and Friess D A 2016 Rates and drivers of mangrove deforestation in Southeast Asia, 2000–2012 *Proc. Natl Acad. Sci.* **113** 344–9
- Robertson A I and Phillips M J 1995 Mangroves as filters of shrimp pond effluent: predictions and biogeochemical research needs *Hydrobiologia* **295** 311–21
- Rockström J, Gaffney O, Rogelj J, Meinshausen M, Nakicenovic N and Schellnhuber H J 2017 A roadmap for rapid decarbonization *Science* **355** 1269–71
- Sanderman J, Hengl T and Fiske G J 2017 Soil carbon debt of 12 000 years of human land use *Proc. Natl Acad. Sci.* **114** 9575–80
- Sanders C J, Maher D T, Tait D R, Williams D, Holloway C, Sippo J Z and Santos I R 2016 Are global mangrove carbon stocks driven by rainfall? *J. Geophys. Res. Biogeosci.* **121** 2600–9
- Siikamäki J, Sanchirico J N and Jardine S L 2012 Global economic potential for reducing carbon dioxide emissions from mangrove loss *Proc. Natl Acad. Sci.* **109** 14369–74
- Simard M, Zhang K, Rivera-Monroy V H, Ross M S, Ruiz P L, Castañeda-Moya E, Twilley R R and Rodriguez E 2006 Mapping height and biomass of mangrove forests in Everglades National Park with SRTM elevation data *Photogramm Eng. Remote Sens.* **72** 299–311
- Snedaker S C 1982 *Mangrove Species Zonation: Why?* (Netherlands: Springer) pp 111–25
- Spalding M, Kainuma M and Collins L 2010 *World Atlas of Mangroves* (London: Earthscan)
- Spalding M, McIvor A, Tonneijck F, Tol S and van Eijk P 2014 *Mangroves for Coastal Defence. Guidelines for Coastal Managers and Policy Makers* (Wetlands International and The Nature Conservancy)
- Stringer C E, Trettin C C and Zarnoch S J 2016 Soil properties of mangroves in contrasting geomorphic settings within the Zambezi River Delta, Mozambique *Wetl. Ecol. Manage.* **24** 139–52
- Tarnocai C, Canadell J G, Schuur E A G, Kuhry P, Mazhitova G and Zimov S 2009 Soil organic carbon pools in the northern circumpolar permafrost region *Glob. Biogeochem. Cycles* **23** GB2023
- R Core Team 2000 *R Language Definition* (Vienna: R Found. Stat. Comput.)
- Temmerman S, Meire P, Bouma T J, Herman P M J, Ysebaert T and De Vriend H J 2013 Ecosystem-based coastal defence in the face of global change *Nature* **504** 79–83
- Valiela I, Bowen J L and York J K 2001 Mangrove forests: one of the world’s threatened major tropical environments *BioScience* **51** 807–15
- Wright M N and Ziegler A 2015 ranger: a fast implementation of random forests for high dimensional data in C++ and R (arXiv:1508.04409)